SKI gene

SKI proto-oncogene

Normal Function

The SKI gene provides instructions for making a protein involved in a signaling pathway that transmits chemical signals from the cell surface to the nucleus. This pathway, called the transforming growth factor beta (TGF- β) pathway, allows the environment outside the cell to affect how the cell produces other proteins. It helps regulate cell growth and division (proliferation), the process by which cells mature to carry out special functions (differentiation), cell movement (motility), and the self-destruction of cells (apoptosis). Through this pathway, a group of proteins called the SMAD complex is turned on (activated). The activated SMAD protein complex moves to the cell nucleus and attaches (binds) to specific areas of DNA to control the activity of particular genes, which help regulate various cellular processes.

The SKI protein controls the activity of the TGF- β pathway by binding to certain SMAD proteins, which interrupts signaling through the pathway. SKI protein binding within the cell can keep the SMAD protein complex from entering the nucleus, so it is unable to activate genes. Binding of the SKI protein can also occur in the nucleus. Although the SMAD complex binds to DNA, the SKI protein attracts other proteins (corepressors) that block its ability to turn genes on.

The SKI protein is found in many cell types throughout the body and appears to play a role in the development of many tissues, including the skull, other bones, skin, and brain.

Health Conditions Related to Genetic Changes


Shprintzen-Goldberg syndrome

At least 10 mutations in the SKI gene have been found in people with Shprintzen-Goldberg syndrome, a condition characterized by distinctive facial features, skeletal abnormalities, and intellectual disability. Most of these mutations change single protein building blocks (amino acids) in the SKI protein. Many of the mutations alter the region of the SKI protein that binds to SMAD proteins. It is thought that altered SKI proteins are unable to attach to SMAD proteins, which allows TGF- β signaling to continue uncontrolled. Excess TGF- β signaling changes the regulation of gene activity and likely disrupts development of many body systems, including the bones and brain, resulting in the wide range of signs and symptoms of Shprintzen-Goldberg syndrome.

Chromosomal Location

Cytogenetic Location: 1p36.33-p36.32, which is the short (p) arm of chromosome 1 between positions 36.33 and 36.32

Molecular Location: base pairs 2,228,695 to 2,310,213 on chromosome 1 (Homo sapiens Annotation Release 108, GRCh38.p7) (NCBI)



Credit: Genome Decoration Page/NCBI

Other Names for This Gene

- proto-oncogene c-Ski
- ski oncogene
- ski oncoprotein
- SKI HUMAN
- v-ski avian sarcoma viral oncogene homolog
- v-ski sarcoma viral oncogene homolog (avian)

Additional Information & Resources

Educational Resources

- Developmental Biology (sixth edition, 2000): The Smad Pathway Activated by TGF-β Superfamily Ligands https://www.ncbi.nlm.nih.gov/books/NBK10043/figure/A1057/
- Genomes (second edition, 2002): SMAD Signaling in Vertebrates https://www.ncbi.nlm.nih.gov/books/NBK21127/box/A7928/#A7929

GeneReviews

 Shprintzen-Goldberg Syndrome https://www.ncbi.nlm.nih.gov/books/NBK1277

Scientific Articles on PubMed

PubMed

https://www.ncbi.nlm.nih.gov/pubmed?term=%28%28SKI%5BTIAB%5D%29+OR+%28v-ski+sarcoma+viral+oncogene+homolog%5BTIAB%5D%29+NOT+%28sphingosine+kinase%29%29+AND+%28%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+720+days%22%5Bdp%5D

OMIM

 V-SKI AVIAN SARCOMA VIRAL ONCOGENE HOMOLOG http://omim.org/entry/164780

Research Resources

- Atlas of Genetics and Cytogenetics in Oncology and Haematology http://atlasgeneticsoncology.org/Genes/GC_SKI.html
- ClinVar https://www.ncbi.nlm.nih.gov/clinvar?term=SKI%5Bgene%5D
- HGNC Gene Family: SKI transcriptional corepressors http://www.genenames.org/cgi-bin/genefamilies/set/748
- HGNC Gene Symbol Report http://www.genenames.org/cgi-bin/gene_symbol_report?q=data/ hgnc_data.php&hgnc_id=10896
- NCBI Gene https://www.ncbi.nlm.nih.gov/gene/6497
- UniProt http://www.uniprot.org/uniprot/P12755

Sources for This Summary

- Carmignac V, Thevenon J, Adès L, Callewaert B, Julia S, Thauvin-Robinet C, Gueneau L, Courcet JB, Lopez E, Holman K, Renard M, Plauchu H, Plessis G, De Backer J, Child A, Arno G, Duplomb L, Callier P, Aral B, Vabres P, Gigot N, Arbustini E, Grasso M, Robinson PN, Goizet C, Baumann C, Di Rocco M, Sanchez Del Pozo J, Huet F, Jondeau G, Collod-Beroud G, Beroud C, Amiel J, Cormier-Daire V, Rivière JB, Boileau C, De Paepe A, Faivre L. In-frame mutations in exon 1 of SKI cause dominant Shprintzen-Goldberg syndrome. Am J Hum Genet. 2012 Nov 2;91(5):950-7. doi: 10.1016/j.ajhg.2012.10.002. Epub 2012 Oct 25.
 - Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/23103230
 Free article on PubMed Central: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3487125/
- Deheuninck J, Luo K. Ski and SnoN, potent negative regulators of TGF-beta signaling. Cell Res. 2009 Jan;19(1):47-57. doi: 10.1038/cr.2008.324. Review.
 Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/19114989
 Free article on PubMed Central: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3103856/

- Suzuki H, Yagi K, Kondo M, Kato M, Miyazono K, Miyazawa K. c-Ski inhibits the TGF-beta signaling pathway through stabilization of inactive Smad complexes on Smad-binding elements. Oncogene. 2004 Jun 24;23(29):5068-76.
 Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/15107821
- OMIM: V-SKI AVIAN SARCOMA VIRAL ONCOGENE HOMOLOG http://omim.org/entry/164780

Reprinted from Genetics Home Reference:

https://ghr.nlm.nih.gov/gene/SKI

Reviewed: February 2013 Published: March 21, 2017

Lister Hill National Center for Biomedical Communications U.S. National Library of Medicine National Institutes of Health Department of Health & Human Services